

### Prognose der lösemittelbedingten Polyneuropathien nach Expositionskarenz

Autoren	Beschwerden			Neurologische Untersuchung			Psychologische Tests			Hirnatrophien			Follow up Jahre
	+	+/-	-	+	+/-	-	+	+/-	--	+	+/-	-	
Bruhn et al. 1981 (n = 26)	6/26	14/26	3/26	0/26	21/26	5/26 <sup>1</sup>	0/26	24/26	2/26	0/26	24/26	2/26	2
Ohrbäk und Lindgren (1988) (n = 30)	5/30 <sup>2</sup>	18/30 <sup>2</sup>	7/30 <sup>2</sup>	8/26	13/26	5/26 <sup>3</sup>	6/23	4/23	13/23	k.A.	k.A.	k.A.	2 - 7
Edlin et al. 1990 (n = 46)	15/46 <sup>2</sup>	11/46 <sup>2</sup>	20/46 <sup>2</sup>	k.A.	k.A.	k.A.	10/46 <sup>4</sup>	16/46 <sup>4</sup>	20/46 <sup>4</sup>	k.A.	k.A.	k.A.	5
Dietz und Triebig 1993 (n=3)	0/3	1/3	2/3	0/3	3/3	0/3	0/3	3/3	0/3	k.A.	k.A.	k.A.	1 - 3

<sup>1</sup> feinschlägiger Tremor, <sup>2</sup> Gedächtnisschwierigkeiten, <sup>3</sup> sensible Polyneuropathie, <sup>4</sup> verbale und logische Testverfahren

Tabelle 2